

16/020281

Welcome to STN International! Enter x:x

LOGINID:sssptaul21bd

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Apr 08 "Ask CAS" for self-help around the clock
NEWS 3 Jun 03 New e-mail delivery for search results now available
NEWS 4 Aug 08 PHARMAMarketLetter(PHARMAML) - new on STN
NEWS 5 Aug 19 Aquatic Toxicity Information Retrieval (AQUIRE)
now available on STN
NEWS 6 Aug 26 Sequence searching in REGISTRY enhanced
NEWS 7 Sep 03 JAPIO has been reloaded and enhanced
NEWS 8 Sep 16 Experimental properties added to the REGISTRY file
NEWS 9 Sep 16 CA Section Thesaurus available in CAPLUS and CA
NEWS 10 Oct 01 CASREACT Enriched with Reactions from 1907 to 1985
NEWS 11 Oct 24 BEILSTEIN adds new search fields
NEWS 12 Oct 24 Nutraceuticals International (NUTRACEUT) now available on STN
NEWS 13 Nov 18 DKILIT has been renamed APOLLIT
NEWS 14 Nov 25 More calculated properties added to REGISTRY
NEWS 15 Dec 04 CSA files on STN
NEWS 16 Dec 17 PCTFULL now covers WP/PCT Applications from 1978 to date
NEWS 17 Dec 17 TOXCENTER enhanced with additional content
NEWS 18 Dec 17 Adis Clinical Trials Insight now available on STN
NEWS 19 Jan 29 Simultaneous left and right truncation added to COMPENDEX,
ENERGY, INSPEC
NEWS 20 Feb 13 CANCERLIT is no longer being updated
NEWS 21 Feb 24 METADEX enhancements
NEWS 22 Feb 24 PCTGEN now available on STN
NEWS 23 Feb 24 TEMA now available on STN
NEWS 24 Feb 26 NTIS now allows simultaneous left and right truncation
NEWS 25 Feb 26 PCTFULL now contains images
NEWS 26 Mar 04 SDI PACKAGE for monthly delivery of multifile SDI results
NEWS 27 Mar 19 APOLLIT offering free connect time in April 2003
NEWS 28 Mar 20 EVENTLINE will be removed from STN
NEWS 29 Mar 24 PATDPAFULL now available on STN
NEWS 30 Mar 24 Additional information for trade-named substances without
structures available in REGISTRY
NEWS 31 Mar 24 Indexing from 1957 to 1966 added to records in CA/CAPLUS
NEWS 32 Apr 11 Display formats in DGENE enhanced
NEWS 33 Apr 14 MEDLINE Reload
NEWS 34 Apr 17 Polymer searching in REGISTRY enhanced

NEWS EXPRESS April 4 CURRENT WINDOWS VERSION IS V6.01a, CURRENT
MACINTOSH VERSION IS V6.0b(ENG) AND V6.0Jb(JP),
AND CURRENT DISCOVER FILE IS DATED 01 APRIL 2003
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer

agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 12:55:54 ON 17 APR 2003

=> fil reg

FILE 'REGISTRY' ENTERED AT 12:56:03 ON 17 APR 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 16 APR 2003 HIGHEST RN 503266-82-8

DICTIONARY FILE UPDATES: 16 APR 2003 HIGHEST RN 503266-82-8

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 6, 2003

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.


Experimental and calculated property data are now available. See HELP PROPERTIES for more information. See STNote 27, Searching Properties in the CAS Registry File, for complete details:
<http://www.cas.org/ONLINE/STN/STNOTES/stnotes27.pdf>

*** YOU HAVE NEW MAIL ***

'REGISTRY' IS DEFAULT FORMAT FOR 'REGISTRY' FILE

=> str


:gra r66,dis



=> d sia

L1 HAS NO ANSWERS

L1 STR



NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 22

STEREO ATTRIBUTES: NONE

=> s l1
 SAMPLE SEARCH INITIATED 13:00:45 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 6 TO ITERATE

100.0% PROCESSED 6 ITERATIONS 5 ANSWERS
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 6 TO 266
 PROJECTED ANSWERS: 5 TO 234

L2 5 SEA SSS SAM L1

=> s l1
 SAMPLE SEARCH INITIATED 13:00:51 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 6 TO ITERATE

100.0% PROCESSED 6 ITERATIONS 5 ANSWERS
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 6 TO 266
 PROJECTED ANSWERS: 5 TO 234

L3 5 SEA SSS SAM L1

=> ful
 FUL IS NOT A RECOGNIZED COMMAND
 The previous command name entered was not recognized by the system.
 For a list of commands available to you in the current file, enter
 "HELP COMMANDS" at an arrow prompt (=>).

=> s 11 ful
FULL SEARCH INITIATED 13:00:58 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 94 TO ITERATE

100.0% PROCESSED 94 ITERATIONS 67 ANSWERS
SEARCH TIME: 00.00.01

L4 67 SEA SSS FUL L1

=> fil caplus
FILE 'CAPLUS' ENTERED AT 13:01:05 ON 17 APR 2003
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 17 Apr 2003 VOL 138 ISS 16
FILE LAST UPDATED: 16 Apr 2003 (20030416/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 14/p
L5 13 L4/P

=> d tot cbib sbs
'SBS' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE
APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data and PI table (default)
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
DMAX ----- MAX, delimited for post-processing
FAM ----- AN, PI and PRAI in table, plus Patent Family data
FBIB ----- AN, BIB, plus Patent FAM
IND ----- Indexing data
IPC ----- International Patent Classifications
MAX ----- ALL, plus Patent FAM, RE
PATS ----- PI, SO
SAM ----- CC, SX, TI, ST, IT
SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;
SCAN must be entered on the same line as the DISPLAY,
e.g., D SCAN or DISPLAY SCAN)
STD ----- BIB, IPC, and NCL

IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels

IMAX ----- MAX, indented with text labels
 ISTD ----- STD, indented with text labels

OBIB ----- AN, plus Bibliographic Data (original)
 OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
 SIBIB ----- IBIB, no citations



HIT ----- Fields containing hit terms
 HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)
 containing hit terms
 HITRN ----- HIT RN and its text modification
 HITSTR ----- HIT RN, its text modification, its CA index name, and
 its structure diagram
 HITSEQ ----- HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and
 its structure diagram
 FHITSEQ ----- First HIT RN, its text modification, its CA index name, its
 structure diagram, plus NTE and SEQ fields
 KWIC ----- Hit term plus 20 words on either side
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.
 ENTER DISPLAY FORMAT (BIB):cbib abs

L5 ANSWER 1 OF 13 CAPLUS COPYRIGHT 2003 ACS
 2003:117704 Document No. 138:172243 Processes for the production of amino-protected derivatives of 4-aminomethylene-pyrrolidin-3-one and/or 4-aminomethylene-pyrrolidin-3-alkoxyimino derivatives and/or gemifloxacin or a salt thereof. Brechtelsbauer, Clemens Michael Helmut; Carpenter, Stephen Thomas; Grinter, Trevor John; Harris, Michael Anthony; Kim, Yeongdae; Kwon, Youngwoon; Lee, Dongchul; Ricard, Francois Xavier; Saunders, Richard Neville (SB Pharmco Puerto Rico Inc., USA; LG Chem Investment Ltd.). PCT Int. Appl. WO 2003011450 A1 20030213, 78 pp.
 DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2.
 APPLICATION: WO 2002-GB3585 20020802. PRIORITY: GB 2001-18938 20010802; GB 2002-17637 20020730.

GI




AB The invention provides a process for the prodn. of a compd. of formula (I): wherein P1 and P2, which may be the same or different, are amino protecting groups, which comprises protection of a compd. of formula (II) in soln. phase continuous operation mode. This confers advantages over batch mode operation. The process is usually conducted in reaction equipment adapted for use in continuous processing mode, for example comprising one or more static mixers or a plug flow reactor. Preferably, the plug flow reactor comprises a jacketed tubular reactor fitted inside with internal mixing elements which continually split and premix the reaction streams promoting mass and heat transfer, whereby a uniform plug flow profile with turbulent fluid flow is achieved. The invention also provides a process for prodn. of the antibacterial compd. gemifloxacin or a pharmaceutically acceptable salt and/or hydrate thereof, comprising converting a compd. of formula (I).

L5 ANSWER 2 OF 13 CAPLUS COPYRIGHT 2003 ACS

2001:693322 Document No. 135:242153 A process for the preparation of pyrrolidino-quinoline-carboxylic acid derivatives (e.g. gemifloxacin) with improved filtration. Kim, Bong Chan (LG Chem Investment Ltd., S. Korea; Kim, Yeong Dae; Choi, Hoon; Kim, Won Sup). PCT Int. Appl. WO 2001068649 A1 20010920, 17 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2001-KR399 20010314. PRIORITY: KR 2000-13011 20000315.

GI




AB The present invention relates to a novel process for prepg. quinoline carboxylic acid antimicrobials I (e.g. gemifloxacin) [Q = CH, CF, CCl, COH, COMe, N; R = H, Me, NH₂; R₁ = cyclopropyl, Et, substituted-Ph; R₂ = H, alkyl, aryl, allyl; R₃, R₄ = H, alkyl, or together with the N atom to which they are attached form a cycle]. For instance, Et₃N, methylcellulose (1.0 wt. % relative to pyrrolidine reactant) and 4-aminomethyl-3-methoxyiminopyrrolidine were added sequentially to an aq. soln. of 3-carboxy-7-chloro-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro[1,8]naphthyridine and allowed to stir at room temp. for 16.5 h. The resulting mixt. was filtered on a glass filter and the product washed. Addn. of the surfactant, methylcellulose, resulted in a faster filtration, 5 min vs. 9 min (scale, 34 - 37 g product). Other process parameters evaluated included sequence of addn. of reagents, variation of reaction temp. and surfactant.

L5 ANSWER 3 OF 13 CAPLUS COPYRIGHT 2003 ACS

2001:380574 Document No. 135:5534 Preparation of quinolonecarboxylates as bactericides and parasiticides. Abdul-Rahman, Shoaab (New Pharma Research

Sweden AB, Swed.). PCT Int. Appl. WO 2001036408 A1 20010525, 123 pp.
 DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2000-SE2217 20001113. PRIORITY: SE 1999-4108 19991115.


GI




AB Title compds. [I; R = NR5R6; R1 = H, (fluoro)alkyl, alkoxyalkyl, etc.; R2,R3 = H, (oxa)alkyl, (di)(alkyl)amino, alkanoyl(oxy), (hetero)aryl, etc.; R4-R6 = groups cited for R2, etc.; NR5R6 = heterocyclyl; R7 = F, Cl, cyano, CO2R1, arylamino, etc.; Z = N or CR9; R9 = groups cited for R2; R4R9 = atoms to complete a ring] were prepd. Thus, 2,4-dichloro-5-fluoroacetophenone was converted in 5 steps to I (R1-R3 = H, R4 = cyclopropyl, R7 = F, Z = CH) (II; R = Cl) which was aminated by 4-picolylamine to give II (R = 4-pyridinylmethylamino). Data for biol. activity of I were given.

L5 ANSWER 4 OF 13 CAPLUS COPYRIGHT 2003 ACS
 2001:185757 Document No. 134:222702 Process for production of naphthyridine-3-carboxylic acid derivatives. Cho, Sungwook; Choi, Hoon; Hayler, John David (Sb Pharmco Puerto Rico Inc., USA; Lg Chemical Ltd.). PCT Int. Appl. WO 2001018002 A1 20010315, 12 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 2000-GB3366 20000901. PRIORITY: GB 1999-20917 19990903.


GI



I



II



III

AB Title compds. I (R = alkyl, haloalkyl) were prepd. by reaction of II (X = a leaving atom/group) with III or its salt. Thus, 5.1 mL Et₃N was added to 3.05 g II (X = Cl) in 25 mL water at 15-20.degree., and the mixt. was stirred for 20 min, after which 3.86 g 4-(aminomethyl)-3-(methoxyimino)pyrrolidinium dimethanesulfonate was added, followed by 5 mL water, and the mixt. was stirred at 20-25.degree. for 17.75 h to give 4.23 g syn-I (R = Me).

L5 ANSWER 5 OF 13 CAPLUS COPYRIGHT 2003 ACS

2001:185719 Document No. 134:222700 Preparation of intermediates for the production of naphthyridinecarboxylic acid-derivative antibiotics.

Grinter, Trevor John; Howie, Simon (SB Pharmco Puerto Rico Inc., USA).




PCT Int. Appl. WO 2001017961 A2 20010315, 9 pp. DESIGNATED STATES: W:

AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG.

(English). CODEN: PIXXD2. APPLICATION: WO 2000-GB3358 20000901.

PRIORITY: GB 1999-20919 19990903.

GI




AB Quinolonecarboxylic acid-deriv. antibiotics (I; R = C1-4 alkyl, C1-4 haloalkyl) are prepd. in high yield and selectivity by the reaction of fluoroquinolonecarboxylic acids (II; X = leaving group) with 4-(aminomethyl)-3-(alkoxyimino)pyrrolidinium dimethanesulfonates (III). Thus, triethylamine was added to 7-chloro-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid, the mixt. stirred and 4-(aminomethyl)-3-(methoxyimino)pyrrolidinium dimethanesulfonate added, producing 7-(3-aminomethyl-4-syn-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid in 86% yield.

L5 ANSWER 6 OF 13 CAPLUS COPYRIGHT 2003 ACS

2000:842133 Document No. 134:17486 Preparation of optically active 7-(pyrrolidin-1-yl)quinolinecarboxylates and -naphthyridinecarboxylates as antibacterials.. Yoon, Sung June; Chung, Yong Ho; Lee, Chi Woo; Lee, Jin Soo; Kim, Nam Doo; Jin, Yoon Ho; Song, Wan Jin; Kim, Ik Hoe; Yang, Wang Yong; Choi, Dong Rack; Shin, Jung Han (Dong Wha Pharm. Ind. Co., Ltd., S. Korea). PCT Int. Appl. WO 2000071541 A1 20001130, 74 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 2000-KR487 20000518. PRIORITY: KR 1999-18158 19990520; KR 2000-24657 20000509.

GI




AB Title compds. (I; Q = CH, CF, CCl, N; Y = H, NH₂; R = alkyl, allyl, PhCH₂), were prepd. Thus, (+)-7-(4-aminomethyl-4-methyl-3-oxopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid hydrochloride (prepn. given) was stirred with methoxylamine hydrochloride in pyridine for 4 h to give 97.5% (-)-7-(4-aminomethyl-4-methyl-3-(Z)-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid hydrochloride. This showed a min. inhibitory concn. of 0.025 .mu.g/mL against Streptococcus pyogenes 308A.

L5 ANSWER 7 OF 13 CAPLUS COPYRIGHT 2003 ACS

1999:576910 Document No. 131:201515 Process for preparing a protected 4-aminomethylpyrrolidin-3-one. Moon, Kwang Yul; Kim, Won Sup; Lee, Tae Hee; Chang, Jay Hyok (LG Chemical Ltd., S. Korea). PCT Int. Appl. WO 9944991 A1 19990910, 30 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 1999-KR99 19990304. PRIORITY: KR 1998-7079 19980304; KR 1998-43636 19981019.

GI



AB Protected 4-aminomethyl-3-pyrrolidinones I (R = CH₂NHP₂, P₁, P₂ = protecting groups) are manufd. by hydrogenating I (R = CN, P₁ = same as in I) in the presence of a Raney-nickel catalyst in a solvent, reacting the resulting aminomethylene deriv. with a compd. to form P₂, and selective redn. of the double bond attached to the ring. This process does not cause formation of an OH group at the 3-position, and the products are useful in the manuf. of quinolone antibiotics. Thus, hydrogenating 20 kg 1-(N-tert-butoxycarbonyl)-4-cyano-3-pyrrolidinone in MeOH-NH₄OH in the presence of a Raney-nickel catalyst, reacting intermediate with Li tert-butoxide in PhMe at .ltoreq.-10.degree., adding di-tert-Bu dicarbonate to complete the reaction, and hydrogenating the 2nd intermediate in PrOH 24 in the presence of Bu₃N and a Pd catalyst gave I (R = NHCOCMe₃, P₂ = COCMe₃) quant.

L5 ANSWER 8 OF 13 CAPLUS COPYRIGHT 2003 ACS


1999:108189 Document No. 130:133551 SB-265805/LB-20304a: fluoronaphthyridinecarboxylic acid antibacterial. Graul, A.; Castaner, J. (Prous Science, Barcelona, 08080, Spain). Drugs of the Future, 23(11), 1199-1204 (English) 1998. CODEN: DRFUD4. ISSN: 0377-8282. Publisher: Prous Science.

AB A review, with refs., of the synthesis, pharmacol., pharmacokinetics and metab. of LB-20304a (a fluoronaphthyridinecarboxylic acid deriv.) (developed by LG Chem.); (later, LB-20304a was given the new code name SB-265805).

L5 ANSWER 9 OF 13 CAPLUS COPYRIGHT 2003 ACS

1999:104541 Document No. 130:168355 Preparation of 7-(4-aminomethyl-3-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid and related compounds as antibacterials.. Hong, Chang Yong; Kim, Young Kwan; Kim, Se Ho; Chang, Jay Hyok; Choi, Hoon; Nam, Do Hyun; Kim, Ae Ri; Lee, Jin Hwa; Park, Ki Sook (LG Chemical Ltd., S. Korea). U.S. US 5869670 A 19990209, 67 pp., Cont. of U.S. Ser. No. 490,978. (English). CODEN: USXXAM. APPLICATION: US 1998-49024 19980327. PRIORITY: KR 1994-13604 19940616; KR 1994-39915 19941230; KR 1994-39930 19941230; US 1995-490978 19950615; US 1997-825992 19970404.

GI




AB Title compds. [I; R = H, Me, amino; Q = CH, CF, CCl, COH, CMe, COMe, N; R1 = cyclopropyl, Et, fluorophenyl; R2 = H, alkyl, cyclopropyl, cyclopropylmethyl, alkynyl, 2-haloethyl, methoxymethyl, methoxycarbonylmethyl, aryl, alkyl, (substituted) PhCH2, pyridylmethyl, etc.; R3, R4 = H, alkyl; R3R4N = ring], were prepd. Thus, 1-cyclopropyl-7-chloro-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid, 4-aminomethylpyrrolidin-3-one O-methyloxime bistrifluoroacetate, and DBU were refluxed in MeCN to give 85% 7-(4-aminomethyl-3-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid. The latter inhibited Staphylococcus aureus 6538p with a min. inhibitory concn. of .1toreq.0.008 .mu.g/mL.

L5 ANSWER 10 OF 13 CAPLUS COPYRIGHT 2003 ACS

1998:677801 Document No. 129:293896 Salt of naphthyridine carboxylic acid derivative.. Kim, Ae Ri; Lee, Jin Hwa; Park, Ki Sook; Choi, Jong Ryoo; Lee, Tae Hee; Chang, Jay Hyok; Nam, Do Hyun; Choi, Hoon (LG Chemical Ltd., S. Korea). PCT Int. Appl. WO 9842705 A1 19981001, 49 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 1998-KR51 19980320. PRIORITY: KR 1997-9840 19970321.

GI



I

AB 7-(3-Aminomethyl-4-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid (I) methanesulfonate and hydrates were prepd., as were pharmaceutical compns. comprising them, and they were used for antibacterial therapy. I methanesulfonate-1.5H₂O and I methanesulfonate-3H₂O were prepd., x-ray diffraction carried out, and stability and antibacterial activity detd.

L5 ANSWER 11 OF 13 CAPLUS COPYRIGHT 2003 ACS
 1998:471466 Document No. 129:122580 Preparation of quinoline(or naphthyridine)-3-carboxylic acids such as 7-(4-aminomethyl-3-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid as antibacterials. Hong, Chang Yong; Kim, Young Kwan; Kim, Se Ho; Chang, Jay Hyok; Choi, Hoon; Nam, Do Hyun; Kim, Ae Ri; Lee, Jin Hwa; Park, Ki Sook (LG Chemical Ltd., S. Korea). U.S. US 5776944 A 19980707, 64 pp., Cont.-in-part of U. S. 5,633,262. (English). CODEN: USXXAM. APPLICATION: US 1997-825992 19970404. PRIORITY: KR 1994-13604 19940616; KR 1994-39915 19941230; KR 1994-39930 19941230; US 1995-490978 19950615.


GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The title compds. [I; R = H, Me, NH₂; Q = CH, CF, CCl, C(OH), C(Me), C(OMe), N; R₁ = cyclopropyl, Et, (un)substituted Ph; R₂ = H, C1-4 alkyl, cyclopropyl, etc.; R₃, R₄ = H, C1-3 alkyl; R₃R₄N = a ring], having an excellent antibacterial activity, were prepd. More specifically, the present invention relates to 7-(4-aminomethyl-3-methoxyiminopyrrolidin-1-yl)-1-cyclopropyl-6-fluoro-4-oxo-1,4-dihydro-1,8-naphthyridine-3-carboxylic acid (II) or its isomers, which was prepd. by reacting a quinolone III [X = halo] with a pyrrolidine oxime IV in the presence of an acid acceptor. (Z)-II isomer has a superior antibacterial activity to the (E)-II isomer (as the free form or as its methanesulfonate) with, e.g., MIC of .ltoreq. 0.008 .mu.g/mL against Staphylococcus aureus 6538p.

L5 ANSWER 12 OF 13 CAPLUS COPYRIGHT 2003 ACS
 1997:638458 Document No. 127:307319 Novel Fluoroquinolone Antibacterial Agents Containing Oxime-Substituted (Aminomethyl)pyrrolidines: Synthesis and Antibacterial Activity of 7-(4-(Aminomethyl)-3-(methoxyimino)pyrrolidin-1-yl)-1-cyclopropyl -6-fluoro-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxylic Acid (LB20304). Hong, Chang Yong; Kim, Young Kwan; Chang, Jay Hyok; Kim, Se Ho; Choi, Hoon; Nam, Do Hyun; Kim, Yong Zu; Kwak, Jin Hwan (Biotech Research Institute, LG Chem Research Park, Tae-Jon, 305-380, S. Korea). Journal of Medicinal Chemistry, 40(22), 3584-3593 (English) 1997. CODEN: JMCMAR. ISSN: 0022-2623. Publisher: American Chemical Society.

GI




AB Title compds. I [X = CF, CCl, CH, COMe, N; R, R1 = H, Me; R2 = Me, Pr, CHMe2, CMe3, CH2Ph, Ph; R3 = H, NH2; R4 = Et, cyclopropyl, 2,4-F2C6H3] were prepd. from the quinolone and the pyrrolidinone fragments. These fluoroquinolones possess potent antimicrobial activity against both Gram-neg. and Gram-pos. organisms, including methicillin-resistant *Staphylococcus aureus* (MRSA). The activity imparted to the substituted quinolone nucleus by the C-8 substituent was in the order F (C5-NH2) > F (C5-H) > naphthyridine > Cl = OMe = H against Gram-pos. organisms. In the case of Gram-neg. strains, activity was in the order F (C5-NH2) > naphthyridine = F (C5-H) > H > Cl > OMe. The advantages provided by the newly introduced oxime group of the quinolones were clearly demonstrated by their comparison to a desoximino compd. In addn., the oxime moiety greatly improved the pharmacokinetic parameters of the novel quinolones. LB20304 (I, X = N, R, R1, R3 = H, R2 = Me, R4 = cyclopropyl) showed the best in vivo efficacy and pharmacokinetic profile in animals, as well as good phys. properties.


L5 ANSWER 13 OF 13 CAPLUS COPYRIGHT 2003 ACS

1996:237481 Document No. 124:289515 Preparation of novel 7-[(4-aminomethyl-3-alkoxyimino)pyrrolidin-1-yl]quinoline-3-carboxylic acid derivatives as antibacterial agents. Kwak, Jin Hwan; Jeong, Yi Na; Oh, Jeong In (LG Chemical Ltd., S. Korea). Eur. Pat. Appl. EP 688772 A1 19951227, 92 pp. DESIGNATED STATES: R: CH, DE, DK, FR, GB, IT, LI, NL, SE. (English). CODEN: EPXXDW. APPLICATION: EP 1995-250143 19950614. PRIORITY: KR 1994-13604 19940616; KR 1994-39915 19941230; KR 1994-39930 19941230.


GI



I



II



III

AB Title compds. I [Q = CH, CF, CCl, C(OH), C(Me), C(OMe), N; R = H, Me, NH₂; R₁ = cyclopropyl, Et, fluoro-substituted phenyl; R₂ = H, C₁-C₄ alkyl, C₃-C₆ alkynyl, (substituted) benzyl, etc.; R₃, R₄ = H, C₁-C₃ alkyl, etc.] were prepd. Reaction of II with III.2CF₃COOH in the presence of 1,8-diazabicyclo[5.4.0]undec-7-ene in MeCN under reflux afforded 85% I [Q = N; R, R₃, R₄ = H; R₁ = cyclopropyl; R₂ = Me] which showed MIC of .1toeq.0.008 .mu.g/mL against Staphylococcus aureus 6538p, Staphylococcus aureus giorgio and Staphylococcus epidermidis 178 vs. 0.25 .mu.g/mL with Ofloxacin.

=> log h

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

33.48

184.64

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-8.46

-8.46

SESSION WILL BE HELD FOR 60 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 13:04:17 ON 17 APR 2003